

The Balance Error Scoring System (BESS)

*Obtain Preseason Baseline Score; Compare with Post-Concussion Score*³³⁻³⁴

The Balance Error Scoring System³³⁻³⁴ provides a portable, cost-effective and objective method of assessing static **postural stability**. The BESS can be used to assess the effects of mild head injury on static postural stability. Information obtained from this clinical balance tool can be used to assist clinicians in making return to play decisions following mild head injury. The BESS can be performed in nearly any environment and takes approximately 10 minutes to conduct.

The balance-testing regime consists three stances on two different surfaces. The three stances are **double leg stance**, **single leg stance** and **tandem stance**. The two different surfaces include both a **firm** (ground) and **foam** surface. **Athletes' stance should consist of the hands on the iliac crests, eyes closed and a consistent foot position depending on the stance.** Shoes should not be worn.

In the **double leg stance**, the feet are flat on the testing surface approximately pelvic width apart.

In the **single leg stance** position, the athlete is to stand on the non-dominant leg with the contralateral limb held in approximately 20° of hip flexion, 45° of knee flexion and neutral position in the frontal plane.

In the **tandem stance** testing position, one foot is placed in front of the other with heel of the anterior foot touching the toe of the posterior foot. The athlete's non-dominant leg is in the posterior position. Leg dominance should be determined by the athlete's kicking preference.

Administering the BESS: Establish baseline score prior to the start of the athletic season. After a concussive injury, re-assess the athlete and compare to baseline score. Only consider return to activity if scores are comparable to baseline score. Use with Standardized Symptom Scale Checklist.

Scoring the BESS: Each of the trials is **20 seconds**. Count the number of errors (deviations) from the proper stance. The examiner should begin counting errors only after the individual has assumed the proper testing position.



Double Leg Stance
Firm Surface



Single Leg Stance
Firm Surface



Tandem Stance
Firm Surface



Double Leg Stance
Foam Surface



Single Leg Stance
Foam Surface



Tandem Stance
Foam Surface

Errors:

- Moving the hands off the hips
- Opening the eyes
- Step, stumble or fall
- Abduction or flexion of the hip beyond 30°
- Lifting the forefoot or heel off of the testing surface
- Remaining out of the proper testing position for greater than 5 seconds

The maximum total number of errors for any single condition is 10.

If a subject commits multiple errors simultaneously, only one error is recorded.

B.E.S.S. SCORECARD

Count Number of Errors max of 10 each stance/surface	FIRM Surface	FOAM Surface
Double Leg Stance (feet together)		
Single Leg Stance (non-dominant foot)		
Tandem Stance (non-dominant foot in back)		
TOTAL SCORES: total each column		
B.E.S.S. TOTAL: (Firm+Foam total)		